

10/729,427

- 2 -

**Amendments to the Claims**

Please amend Claims 1 and 14 to correct typographical errors.

Please cancel Claims 25-55. Applicants reserve the right to file a continuing application or take such other appropriate action as deemed necessary to protect the non-elected inventions. Applicants do not hereby abandon or waive any rights in the non-elected inventions.

Rejoinder of Claims 5-9, 16-19 and 24, withdrawn in response to the requirement for the election of species, is hereby requested.

The Claim Listing below will replace all prior versions of the claims in the application:

**Claim Listing**

1. (Currently Amended) A method of treating a patient suffering from an inflammatory condition, comprising  
treating said patient with a therapeutically effective amount of a cholinergic agonist selective for an  $\alpha 7$  nicotinic receptor,  
wherein said condition is selected from the group consisting of appendicitis, peptic, gastric and duodenal ulcers, peritonitis, pancreatitis, epiglottitis, achalasia, cholangitis, cholecystitis, hepatitis, Whipple's disease, asthma, allergy, anaphylactic shock, immune complex disease, organ ischemia, reperfusion injury, organ necrosis, hay fever, sepsis, septicemia, endotoxic shock, cachexia, hyperpyrexia, eosinophilic granuloma, granulomatosis, sarcoidosis, septic abortion, epididymitis, vaginitis, prostatitis, urethritis, bronchitis, emphysema, rhinitis, cystic fibrosis, pneumonitis, pneumoultramicroscopic silicovolcanoconiosis, alveolitis, bronchiolitis, pharyngitis, pleurisy, sinusitis, influenza, respiratory syncytial virus infection, herpes infection, HIV infection, hepatitis B virus infection, hepatitis C virus infection, disseminated bacteremia, Dengue fever, candidiasis, malaria, filariasis, amebiasis, hydatid cysts, burns, vasculitis, angitis, endocarditis, arteritis, atherosclerosis, thrombophlebitis, pericarditis, myocarditis, myocardial ischemia, periarteritis nodosa, rheumatic fever, coeliac disease, congestive heart failure, adult respiratory distress syndrome, chronic obstructive pulmonary disease, meningitis, encephalitis, neuritis, neuralgia, spinal cord injury,


10/729,427

- 3 -

paralysis, uveitis, arthritides, arthralgias, osteomyelitis, fasciitis, Paget's disease, gout, periodontal disease, rheumatoid arthritis, synovitis, myasthenia gravis, thyroiditis, systemic lupus erythematosus, Goodpasture's syndrome, Behcets's syndrome, allograft rejection, graft-versus-host disease, ~~ankylosing spondylitis, Berger's disease,~~ ankylosing spondylitis, Berger's disease, Retier's syndrome, and Hodgkins disease.

2-3. (Cancelled)


4. (Original) The method of claim 1, wherein the cholinergic agonist is selected from the group consisting of a quaternary analog of cocaine; (1-aza-bicyclo[2.2.2]oct-3-yl)-carbamic acid 1-(2-fluorophenyl)-ethyl ester; a compound of formula I:



wherein, R represents hydrogen or methyl, and  
n represents 0 or 1; a pharmaceutically acceptable salt of a compound of formula I; a compound of formula II:

10/729,427

- 4 -



wherein:

m is 1 or 2,

n is 0 or 1,

Y is CH, N or NO,

X is oxygen or sulfur,

W is oxygen, H<sub>2</sub> or F<sub>2</sub>,

A is N or C(R<sup>2</sup>),

G is N or C(R<sup>3</sup>),

D is N or C(R<sup>4</sup>),

with the proviso that no more than one of A, G and D is nitrogen but at least one of Y, A, G and D is nitrogen or NO,

R<sup>1</sup> is hydrogen or C<sub>1</sub>-C<sub>4</sub> alkyl,

R<sup>2</sup>, R<sup>3</sup> and R<sup>4</sup> are independently hydrogen, halogen, C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>2</sub>-C<sub>4</sub> alkenyl,

C<sub>2</sub>-C<sub>4</sub> alkynyl, aryl, heteroaryl, OH, OC<sub>1</sub>-C<sub>4</sub> alkyl, CO<sub>2</sub>R<sup>1</sup>, -CN, -NO<sub>2</sub>, -NR<sup>5</sup>R<sup>6</sup>, -CF<sub>3</sub> or -

OSO<sub>2</sub>CF<sub>3</sub>, or R<sup>2</sup> and R<sup>3</sup>, R<sup>3</sup> and R<sup>4</sup>, respectively, may together form another six

membered aromatic or heteroaromatic ring sharing A and G, or G and D, respectively,

containing between zero and two nitrogen atoms, and substituted with one to two of the

following substituents: independently hydrogen, halogen, C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>2</sub>-C<sub>4</sub> alkenyl, C<sub>2</sub>-

C<sub>4</sub> alkynyl, aryl, heteroaryl, OH, OC<sub>1</sub>-C<sub>4</sub> alkyl, CO<sub>2</sub>R<sup>1</sup>, -CN, -NO<sub>2</sub>, -NR<sup>5</sup>R<sup>6</sup>, -CF<sub>3</sub> or -

OSO<sub>2</sub>CF<sub>3</sub>,

10/729,427

- 5 -


$R^5$  and  $R^6$  are independently hydrogen,  $C_1$ - $C_4$  alkyl,  $C(O)R^7$ ,  $C(O)NHR^8$ ,  $C(O)OR^9$ ,  $SO_2R^{10}$  or may together be  $(CH_2)_jQ(CH_2)_k$  where  $Q$  is O, S,  $NR^{11}$ ,

or a bond,

$j$  is 2 to 7,


$k$  is 0 to 2,

$R^7$ ,  $R^8$ ,  $R^9$ ,  $R^{10}$  and  $R^{11}$  are independently  $C_1$ - $C_4$  alkyl, aryl, or heteroaryl, or an enantiomer thereof; a pharmaceutically acceptable salt of a compound of formula II; a compound of formula III:



III

wherein  $R_1$ ,  $R_6$  and  $R_7$  are hydrogen or  $C_1$ - $C_4$  alkyl, and  $R_2$  is selected from a group of




and

10/729,427


- 6 -

wherein,  $R_3$ ,  $R_4$  and  $R_5$  are selected from the group consisting of hydrogen,  $C_1$ - $C_4$  alkyl optionally substituted with N,N-dialkylamino having 1 to 4 carbons in each of the alkyls,  $C_1$ - $C_6$  alkoxy optionally substituted with N,N-dialkylamino having 1 to 4 carbons in each of the alkyls, carboalkoxy having 1 to 4 carbons in the alkoxy, amino, amido having 1 to 4 carbons in the acyl, cyano, and N,N-dialkylamino having 1 to 4 carbons in each of the alkyls, halo, hydroxyl or nitro; and a compound of formula IV:



wherein X is O or S, and R is selected from the group consisting of H,  $OR_1$ ,  $NHC(O)R_1$ , and a halogen, wherein  $R_1$  is a  $C_1$ - $C_4$  alkyl.

5. (Original) The method of claim 1, wherein the cholinergic agonist is a compound of formula I:




wherein, R represents hydrogen or methyl, and  
n represents 0 or 1;  
or a pharmaceutically acceptable salt thereof.


10/729,427

- 7 -

6. (Original) The method of claim 5, wherein the cholinergic agonist is (-)-spiro[1-azabicyclo[2.2.2]octane-3,5'-oxazolidin-2'-one]



7. (Original) The method of claim 1, wherein the cholinergic agonist is a compound of formula II:



wherein:

$m$  is 1 or 2;

$n$  is 0 or 1;

Y is CH, N or NO;

X is oxygen or sulfur;

W is oxygen,  $H_2$  or  $F_2$ ;

$A$  is  $\mathbb{N}$  or  $C(\mathbb{R}^2)$ ;

10/729,427

- 8 -

G is N or C(R<sup>3</sup>);

D is N or C(R<sup>4</sup>);

with the proviso that no more than one of A, G and D is nitrogen but at least one of Y, A, G and D is nitrogen or NO;

R<sup>1</sup> is hydrogen or C<sub>1</sub>-C<sub>4</sub> alkyl;

R<sup>2</sup>, R<sup>3</sup> and R<sup>4</sup> are independently hydrogen, halogen, C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>2</sub>-C<sub>4</sub> alkenyl, C<sub>2</sub>-C<sub>4</sub> alkynyl, aryl, heteroaryl, OH, OC<sub>1</sub>-C<sub>4</sub> alkyl, CO<sub>2</sub>R<sup>1</sup>, -CN, -NO<sub>2</sub>, -NR<sup>5</sup>R<sup>6</sup>, -CF<sub>3</sub> or -OSO<sub>2</sub>CF<sub>3</sub>, or R<sup>2</sup> and R<sup>3</sup>, R<sup>3</sup> and R<sup>4</sup>, respectively, may together form another six membered aromatic or heteroaromatic ring sharing A and G, or G and D, respectively, containing between zero and two nitrogen atoms, and substituted with one to two of the following substituents: independently hydrogen, halogen, C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>2</sub>-C<sub>4</sub> alkenyl, C<sub>2</sub>-C<sub>4</sub> alkynyl, aryl, heteroaryl, OH, OC<sub>1</sub>-C<sub>4</sub> alkyl, CO<sub>2</sub>R<sup>1</sup>, -CN, -NO<sub>2</sub>, -NR<sup>5</sup>R<sup>6</sup>, -CF<sub>3</sub> or -OSO<sub>2</sub>CF<sub>3</sub>;

R<sup>5</sup> and R<sup>6</sup> are independently hydrogen, C<sub>1</sub>-C<sub>4</sub> alkyl, C(O)R<sup>7</sup>, C(O)NHR<sup>8</sup>, C(O)OR<sup>9</sup>, SO<sub>2</sub>R<sup>10</sup> or may together be (CH<sub>2</sub>)<sub>j</sub>Q(CH<sub>2</sub>)<sub>k</sub> where Q is O, S, NR<sup>11</sup>, or a bond;

j is 2 to 7;

k is 0 to 2;


R<sup>7</sup>, R<sup>8</sup>, R<sup>9</sup>, R<sup>10</sup> and R<sup>11</sup> are independently C<sub>1</sub>-C<sub>4</sub> alkyl, aryl, or heteroaryl, or an enantiomer thereof, or a pharmaceutically acceptable salts thereof.

8. (Original) The method of claim 7, wherein the cholinergic agonist is a compound of formula II wherein m is 1; n is 0; p is 0; x is oxygen; A is C(R<sup>2</sup>); G is C(R<sup>3</sup>); and D is C(R<sup>4</sup>).
9. (Original) The method of claim 7, wherein the cholinergic agonist is 5'-phenylspiro[1-azabicyclo[2.2.2]octane-3,2'-(3'H)-furo[2,3-b]pyridin].

10/729,427


- 9 -

10. (Original) The method of claim 1, wherein the cholinergic agonist is a compound of formula III:



(III)

wherein  $R_1$ ,  $R_6$  and  $R_7$  are hydrogen or  $C_1$ - $C_4$  alkyl; and  $R_2$  is selected from a group of



and wherein,  $R_3$ ,  $R_4$  and  $R_5$  are selected from the group consisting of hydrogen,  $C_1$ - $C_4$  alkyl optionally substituted with N,N-dialkylamino having 1 to 4 carbons in each of the alkyls,  $C_1$ - $C_6$  alkoxy optionally substituted with N,N-dialkylamino having 1 to 4 carbons in each of the alkyls, carboalkoxy having 1 to 4 carbons in the alkoxy, amino, amido having 1 to 4 carbons in the acyl, cyano, and N,N-dialkylamino having 1 to 4 carbons in each of the alkyls, halo, hydroxyl or nitro.




10/729,427

- 10 -


11. (Original) The method of claim 10, wherein the cholinergic agonist is a compound of formula III, wherein R<sub>2</sub> is attached to the 3-position of the tetrahydropyridine ring, and further wherein R<sub>3</sub>, which is attached to the 4- or the 2- position of the phenyl ring, is selected from the group consisting of amino, hydroxyl, chloro, cyano, dimethylamino, methyl, methoxy, acetylamino, acetoxy, and nitro.
12. (Original) The method of claim 10, wherein the cholinergic agonist is a compound selected from the group consisting of formula III, wherein R<sub>3</sub> is hydroxyl, and wherein R<sub>1</sub>, R<sub>4</sub>, and R<sub>5</sub> are hydrogen; formula III, wherein R<sub>3</sub> is acetylamino and wherein R<sub>1</sub>, R<sub>4</sub>, and R<sub>5</sub> are hydrogen; formula III, wherein R<sub>3</sub> is acetoxy and wherein R<sub>1</sub>, R<sub>4</sub>, and R<sub>5</sub> are hydrogen; formula III, wherein R<sub>3</sub> is methoxy, and wherein R<sub>1</sub>, R<sub>4</sub>, and R<sub>5</sub> are hydrogen; formula III, wherein R<sub>3</sub> is methoxy and wherein R<sub>1</sub> and R<sub>4</sub> are hydrogen, and further wherein R<sub>3</sub> is attached to the 2-position of the phenyl ring, and R<sub>5</sub>, which is attached to the 4-position of the phenyl ring, is methoxy or hydroxy.
13. (Original) The method of claim 10, wherein the cholinergic agonist is selected from the group consisting of 3-2,4-dimethoxybenzylidene anabaseine (DMXB-A), 3-(4-hydroxybenzylidene)anabaseine, 3-(4-methoxybenzylidene)anabaseine, 3-(4-aminobenzylidene)anabaseine, 3-(4-hydroxy-2-methoxybenzylidene)anabaseine, 3-(4-methoxy-2-hydroxybenzylidene)anabaseine, trans-3-cinnamylidene anabaseine, trans-3-(2-methoxy-cinnamylidene)anabaseine and trans-3-(4-methoxycinnamylidene)anabaseine.
14. (Currently Amended) The method of claim 10, wherein the cholinergic agonist is 3-(4-hydroxy-2-methoxybenzylidene) ~~anabaseine~~ anabaseine

10/729,427


- 11 -



15. (Original) The method of claim 10, wherein the cholinergic agonist is 3-(2,4-dimethoxybenzylidene)anabaseine.



16. (Original) The method of claim 1, wherein the cholinergic agonist is a compound of formula IV:



wherein X is O or S; and

R is selected from the group consisting of H, OR<sub>1</sub>, NHC(O)R<sub>1</sub>, and a halogen, wherein R<sub>1</sub> is a C<sub>1</sub>-C<sub>4</sub> alkyl.

10/729,427

- 12 -

17. (Original) The method of claim 15, wherein the cholinergic agonist is selected from a group consisting of N-[(3R)-1-azabicyclo[2.2.2]oct-3-yl]-4-(4-hydroxyphenoxy)benzamide, N-[(3R)-1-azabicyclo[2.2.2]oct-3-yl]-4-(4-acetamidophenoxy)benzamide, N-[(3R)-1-azabicyclo[2.2.2]oct-3-yl]-4-(phenylsulfonyl)benzamide, and N-[(3R)-1-azabicyclo[2.2.2]oct-3-yl]-4-(3-chlorophenylsulfonyl)benzamide.
18. (Original) The method of claim 15, wherein the cholinergic agonist is N-[(3R)-1-azabicyclo[2.2.2]oct-3-yl]-4-(phenylsulfonyl)benzamide.
19. (Original) The method of claim 1, wherein the cholinergic agonist is cocaine methiodide.
20. (Original) The method of claim 1 wherein the condition is selected from the group consisting of appendicitis, peptic, gastric and duodenal ulcers, peritonitis, pancreatitis, hepatitis, asthma, allergy, anaphylactic shock, organ necrosis, hay fever, sepsis, septicemia, endotoxic shock, cachexia, septic abortion, disseminated bacteremia, burns, coeliac disease, congestive heart failure, adult respiratory distress syndrome, chronic obstructive pulmonary disease, rheumatoid arthritis, systemic lupus erythematosus, myocardial ischemia, spinal cord injury, paralysis, allograft rejection and graft-versus-host disease.
21. (Original) The method of claim 1 wherein the condition selected from the group consisting of appendicitis, peptic, gastric or duodenal ulcers, peritonitis, pancreatitis, hepatitis, asthma, allergy, anaphylactic shock, organ necrosis, hay fever, sepsis, septicemia, endotoxic shock, cachexia, septic abortion, disseminated bacteremia, burns, congestive heart failure, adult respiratory distress syndrome, chronic obstructive pulmonary disease, rheumatoid arthritis, systemic lupus erythematosus, myocardial ischemia, cerebral infarction, cerebral embolism, spinal cord injury, paralysis, allograft rejection or graft-versus-host disease.
22. (Original) The method of Claim 1 wherein the condition is selected from the group consisting of peritonitis, pancreatitis, sepsis, endotoxic shock, adult respiratory distress

10/729,427

- 13 -

syndrome, chronic obstructive pulmonary disease, rheumatoid arthritis, systemic lupus erythematosus, myocardial ischemia, allograft rejection, asthma, graft-versus-host-disease, congestive heart failure and cystic fibrosis.

23. (Original) The method of claim 1, wherein the condition selected from the group consisting of peritonitis, pancreatitis, sepsis, endotoxic shock, cachexia, adult respiratory distress syndrome, chronic obstructive pulmonary disease, rheumatoid arthritis, systemic lupus erythematosus, myocardial ischemia, and allograft rejection.
24. (Original) The method of claim 1, wherein the condition is sepsis.
- 25-55. (Cancelled)

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

☒ **BLACK BORDERS**

☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**

☐ **FADED TEXT OR DRAWING**

☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**

☐ **SKEWED/SLANTED IMAGES**

☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**

☐ **GRAY SCALE DOCUMENTS**

☒ **LINES OR MARKS ON ORIGINAL DOCUMENT**

☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**

☒ **OTHER:** \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**